arXiv:cond-mat/0104188v1 [cond-mat.stat-mech] 11 Apr 2001

Photon Counting Statistics for Single Molecule Spectroscopy

Eli Barkai, YounJoon Jung, and Robert Silbey
Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139.

(February 1, 2008)

Abstract

We present a general relation between Mandel's Q parameter describing variance of number of photons emitted from a single molecule and a three time correlation function describing a spectral diffusion process the molecule is undergoing. For a prototype of such a stochastic process an exact solution is found which exhibits rich types of physical behaviors.

Pacs: 78., 42.50.Ar, 33.80.-b, 05.40.-a

Experimental advances have made it possible to measure the spectral line of a single molecule (SM) embedded in a condensed phase [1]. The statistical properties of the photons emitted from such a SM illuminates the interplay between dynamical processes in the condensed phase, SM dynamics and light-matter interaction. An important process responsible for the emission pattern of a SM is spectral diffusion, i.e., perturbers such as two level systems or excitations in the environment of the excited SM lead to random changes in the transition frequency of the SM and in this way the SM comes in and out of resonance with the exciting laser field [2–11]. The fundamental issue addressed in this work is the photon statistics of a single molecule undergoing a spectral diffusion process. Photon statistics of an ensemble of molecules will clearly differ from that of a SM due to inhomogeneous broadening which is absent in SM spectroscopy (SMS) and due to correlation between fluorescence photons which exists only on the SM level.

We will model these fluctuations in a semiclassical way, using the Bloch equation, in the limit of weak laser field. The spectral diffusion process is described using a Kubo-Anderson sudden jump approach [12]. The same model leads to a basic understanding of line shape phenomena, namely of the average number of counts $\langle n \rangle$ per measurement time T. Variants of this model have been used in the past fifty years to model ensemble measurements in e.g., NMR [12] and non-linear spectroscopy [13]. More recently, this approach was applied to model SMS in glass, by Geva and Skinner [14] and in [11] to describe static properties of line shapes and by Plakhotnik [10] to model time dependent fluctuations of SMS.

Here we present a general formula for Mandel's Q parameter,

$$Q = \frac{\langle n^2 \rangle - \langle n \rangle^2}{\langle n \rangle} - 1$$

where n is the random number of counts. We show how Q is related to the underlying stochastic events and how Q yields new information not contained in the average $\langle n \rangle$. For a simple Kubo-Anderson spectral diffusion process we find an exact solution for Q which gives insight on characteristic behaviors of SMS fluctuations. If $Q = 0$ the photon count is Poissonian while our semiclassical results show a super Poissonian behavior $Q > 0$. For short times (e.g., ns), the correlation function between fluorescence photons emitted by a single molecule shows anti-bunching, a sub-Poissonian non-classical effect [15,16]. Our approach works well for longer times such that $T\Gamma \gg 1$ where Γ is the radiative decay rate of the molecule.

The key quantity in the present formulation is a three time correlation function $\mathrm{C}_3(\tau_1, \tau_2, \tau_3)$, which is similar to non-linear response functions investigated in the context of four wave mixing processes [13]. The three time correlation function contains all the microscopic information relevant for the calculation of Q. It appeared previously in a recent work of Plakhotnik [10] in the context of intensity–time–frequency–correlation technique. In this letter important time ordering properties of this function are investigated, and Q is evaluated. The relation between $\mathrm{C}_3(\tau_1, \tau_2, \tau_3)$ and line shape fluctuations described by Q generalizes the Wiener-Khintchine theorem, which gives the relation between the averaged line shape and the one time dipole correlation function.

An important issue is the fast and slow modulation limits, to be defined later. If the bath is fast the line exhibits the well known behavior of motional narrowing (as the time scale of the bath becomes short; the line is narrowed). Considering a simple spectral diffusion process, we show that Q also exhibits motional narrowing, though in this limit the fluctuations described by Q become small and when time scale of the bath goes to zero we find Poissonian statistics. If the bath is slow, Q becomes large, and a simple adiabatic type of approximation, based on steady state solution of time independent Bloch equation, holds. Our exact results are evaluated for an arbitrary time scale of the bath and are shown to interpolate between the fast and slow modulation limits. We note that several authors have analyzed intensity fluctuations in SMS using the slow modulation assumption [3–5,10], here we extend these results to include the physically important case of fast modulation.

We assume a simple non-degenerate two state single molecule in an external classical laser field. The first electronic excited state $|e\rangle$ is located at energy ω_0 above the ground state $|g\rangle$. We consider the SM Hamiltonian

$$H = \frac{\hbar}{2}\omega_0\sigma_z + \sum_{j=1}^{J}\frac{\hbar}{2}\Delta\omega_j(t)\sigma_z - \mathbf{d}\cdot\mathbf{E}\cos\left(\omega_{\rm L}t\right), \quad (1)$$

where σ_z is the Pauli matrix. The second term in H reflects the effect of the time evolution of the environment on the absorption frequency of the SM coupled to J perturbers. The last term in Eq. (1) describes the interaction between the SM and the laser field (frequency $\omega_{\rm L}$), while $\mathbf{d} \equiv \mathbf{d}_{\rm eg}\sigma_x$ is the dipole operator with the real matrix element $\mathbf{d}_{\rm eg} = \langle e|\mathbf{d}|g\rangle$.

The molecule is described by 2×2 density matrix ρ whose elements are $\rho_{\rm ee}, \rho_{\rm gg}, \rho_{\rm eg}, \rho_{\rm ge}$. Let, $u = (\rho_{\rm ge}e^{-i\omega_{\rm L}t} + \rho_{\rm eg}e^{i\omega_{\rm L}t})/2$, $v = -i(\rho_{\rm ge}e^{-i\omega_{\rm L}t} - \rho_{\rm eg}e^{i\omega_{\rm L}t})/2$, and $w = (\rho_{\rm ee} - \rho_{gg})/2$. Using Eq. (1) the stochastic Bloch equations, in the rotating wave approximation, are

$$\begin{aligned}\dot{u} &= \delta_{\rm L}(t)v - \Gamma u/2,\\ \dot{v} &= -\delta_{\rm L}(t)u - \Omega w - \Gamma v/2,\\ \dot{w} &= \Omega v - \Gamma w - \Gamma/2,\end{aligned} \quad (2)$$

$\Omega = -\mathbf{d}_{\rm eg}\cdot\mathbf{E}/\hbar$ is the Rabi frequency, $\delta_{\rm L}(t) = \omega_{\rm L} - \omega_0 - \sum_{j=1}^{J}\Delta\omega_j(t)$ is the detuning frequency and without loss of generality we set $\omega_0 = 0$. SM experiments [17,18] were shown to be compatible with the *deterministic* two level optical Bloch equation approach and this gives further justification for our assumptions.

According to semiclassical theory of photon counting statistics [19], the probability of recording n photons in time interval $(0, T)$ is given by

$$p(n, T) = \left\langle \frac{W^n}{n!}\exp\left(-W\right)\right\rangle \quad (3)$$

with $W = \xi\int_0^T I(t)\mathrm{d}t$, where $I(t)$ is the photon current and ξ is a suitable constant depending on detection efficiency. The semiclassical approach is valid for large number of counts, when the stream of incoming absorption photons (i.e., Ωv) is equal to the stream of emitted fluorescence photons (i.e., $\Gamma\rho_{\rm ee}$). We use $W \equiv \xi\Omega\int_0^T v(t)\mathrm{d}t$, which is ξ times the work of the driving field per unit energy defined by $\hbar\omega_{\rm L}$, using the Bloch equation it is easy to show that when $W >> 1$ (i.e., photon count is large) $W \to \xi\Gamma\int_0^T \rho_{\rm ee}(t)\mathrm{d}t$ as expected. Using Eq. (3) the average number of photons counted in time interval $(0, T)$ is $\langle n\rangle = \langle W\rangle$ where $\langle\cdots\rangle$ is an average over the stochastic process. Mandel's Q parameter is used to characterize the fluctuations and it is straightforward to show that

$$Q = \frac{\langle W^2\rangle - \langle W\rangle^2}{\langle W\rangle}. \quad (4)$$

We see that $Q \geq 0$ indicating that photon statistics is super Poissonian.

We now consider the important limit of weak laser intensity of Eq. (2). We use a straightforward perturbation expansion in the Rabi frequency to find

$$v = \frac{\Omega}{2}\mathrm{Re}\left\{\int_0^t \mathrm{d}t_1 \exp\left[-i\int_{t_1}^{t}\mathrm{d}t'\delta_{\rm L}(t') - \Gamma\frac{(t-t_1)}{2}\right]\right\}. \quad (5)$$

In standard line shape theories Eq. (5) is averaged, the well known result gives the Wiener–Khintchine formula for the line shape $\langle I(\omega_{\rm L})\rangle = \lim_{T\to\infty}\langle W\rangle/T$, which is valid for stationary processes

$$\langle I\left(\omega_{\rm L}\right)\rangle = \frac{\xi\Omega^2}{2}\mathrm{Re}\left[\int_0^\infty \mathrm{d}\tau e^{-i\omega_{\rm L}\tau - \Gamma\tau/2}\mathrm{C}_1\left(\tau\right)\right], \quad (6)$$

where the one time dipole correlation function is $\mathrm{C}_1(\tau) = \langle e^{i\int_0^\tau \sum_j \Delta_j\omega(t')\mathrm{d}t'}\rangle$. Let us now consider the fluctuation,

$$\langle W^2\rangle = \frac{\xi^2\Omega^4}{16}\int_0^T\int_0^T\int_0^T\int_0^T \mathrm{d}t_1\mathrm{d}t_2\mathrm{d}t_3\mathrm{d}t_4\times$$
$$e^{-i\omega_{\rm L}(t_2-t_1+t_3-t_4)-\Gamma(|t_1-t_2|+|t_3-t_4|)/2}\mathrm{C}_3\left(\tau_1,\tau_2,\tau_3\right) \quad (7)$$

where the three time correlation function

$$\mathrm{C}_3\left(\tau_1,\tau_2,\tau_3\right) =$$
$$\langle\exp\left[i\int_{t_1}^{t_2}\sum_j\Delta\omega_j(t')\mathrm{d}t' - i\int_{t_3}^{t_4}\sum_j\Delta\omega_j(t')\mathrm{d}t'\right]\rangle, \quad (8)$$

contains the information on the *stationary* spectral diffusion process relevant for the calculation of Q. It depends on the time ordering of t_1, t_2, t_3, t_4 for which there are $4! = 24$ options. In Eq. (8) $\tau_1 = t_{\rm II} - t_{\rm I}$, $\tau_2 = t_{\rm III} - t_{\rm II}$ and $\tau_3 = t_{\rm IV} - t_{\rm III}$ and by definition $t_{\rm I} < t_{\rm II} < t_{\rm III} < t_{\rm IV}$. Let us consider the case $t_1 < t_2 < t_3 < t_4$ (for this case $t_1 = t_{\rm I}, t_2 = t_{\rm II}$ etc) and $\tau_1 = t_2 - t_1$, $\tau_2 = t_3 - t_2$ and $\tau_3 = t_4 - t_3$. We define the pulse function $S(t) = -1$ for $t_{\rm I} < t < t_{\rm II}$, $S(t) = 0$ for $t_{\rm II} < t < t_{\rm III}$ and $S(t) = 1$ when $t_{\rm III} < t < t_{\rm IV}$, the shape of this pulse is shown in the first line of Table 1. Using this pulse the correlation function is written as a characteristic functional, $\mathrm{C}_3\left(\tau_1,\tau_2,\tau_3\right) = \langle\exp\left[-i\int_{t_{\rm I}}^{t_{\rm IV}} S(t')\sum\Delta\omega_j(t')\mathrm{d}t'\right]\rangle$. Similarly other pulse shapes describe the other time orderings as described in Table 1. We note that pulses 1 and 2 are encountered in theory of four wave mixing [13].

Eqs. (4-8) give a general prescription for the calculation of Q for broad types of spectral diffusions. Now we investigate basic properties of line shape fluctuations considering a simple process [12]. We assume $J = 1$ and $\Delta\omega(t) = \nu h(t)$ and $h(t)$ is a random telegraph process $h(t) = 1$ or $h(t) = -1$. The transition rate from state up to down and vice versa is R, generalizations for more complicated cases will be considered elsewhere.

Using a method of Suárez and Silbey [20], developed in the context of photon echo experiments, we now analyze the properties of the three time correlation function. Define the weights $P^s_{if}(t) = \langle\exp\left[-is\int_0^t\Delta\omega(t')\mathrm{d}t'\right]\rangle_{if}$

where the initial (final) state of the stochastic process $\Delta\omega$ is i (f) and $s = 0$ or $s = \pm 1$ or $s = \pm 2$ [21]. For example $P_{++}^{-1}(t)$ is the value of $\langle \exp\left(i\int_0^t \Delta\omega(t')\mathrm{d}t'\right)\rangle$ for a path restricted to have $\Delta\omega(0) = \nu$ and $\Delta\omega(t) = \nu$. Using these weights and based on the Markovian property of the process, we find for $t_1 < t_2 < t_3 < t_4$

$$\mathrm{C}_3\left(\tau_1,\tau_2,\tau_3\right) = \frac{1}{2}\sum_{i,j,k,l} P_{ij}^{-1}\left(\tau_1\right)P_{jk}^{0}\left(\tau_2\right)P_{kl}^{1}\left(\tau_3\right)$$

where the summations are over all possible values of $i = \pm$, $j = \pm$,$k = \pm$, and $l = \pm$.

	$S(t)$	$\mathrm{C}_3(\tau_1,\tau_2,\tau_3)$	time order
1	 1 0 -1	$\frac{1}{2}\sum P_{ij}^{-1}(\tau_1)P_{jk}^{0}(\tau_2)P_{kl}^{1}(\tau_3)$	$1234_{1\to 4,2\to 3}$
2	 1 0	$\frac{1}{2}\sum P_{ij}^{1}(\tau_1)P_{jk}^{0}(\tau_2)P_{kl}^{1}(\tau_3)$	$2134_{1\to 4,2\to 3}$
3	 2 1	$\frac{1}{2}\sum P_{ij}^{1}(\tau_1)P_{jk}^{2}(\tau_2)P_{kl}^{1}(\tau_3)$	$2314_{1\to 4,2\to 3}$

Table 1. Each pulse in column 1 represents four time orderings corresponding to a three time correlation function in column 2. The symbol $1234_{1\to 4,2\to 3}$ corresponds to $t_1 < t_2 < t_3 < t_4$, $t_4 < t_2 < t_3 < t_1$, $t_1 < t_3 < t_2 < t_4$ and $t_4 < t_3 < t_2 < t_1$. The complex conjugate (C.C) of $\mathrm{C}_3\left(\tau_1,\tau_2,\tau_3\right)$ in line 1 describes the time orderings $2143_{1\to 4,2\to 3}$, C.C $[\mathrm{C}_3\left(\tau_1,\tau_2,\tau_3\right)]$ of pulse 2 corresponds to $1243_{1\to 4,2\to 3}$, while C.C $[\mathrm{C}_3\left(\tau_1,\tau_2,\tau_3\right)]$ of pulse 3 corresponds to $1423_{1\to 4,2\to 3}$.

In a similar way we consider the other 23 time orderings. We find only three types of pulses $S(t)$ are needed for the definition of the three time correlation function, these pulses are presented in Table 1 together with the corresponding $\mathrm{C}_3(\tau_1,\tau_2,\tau_3)$. We then sum all the 24 contributions and use the convolution theorem of Laplace transform to arrive at the main result of this paper

$$\begin{aligned}\langle W^2\rangle = \mathcal{L}^{-1}\{ & \frac{\Omega^4\xi^2}{16u^2}\sum_{i,j,k,l}[\\ & \hat{P}_{ij}^{-1}\left(u+u_+\right)\hat{P}_{jk}^{0}\left(u\right)\hat{P}_{kl}^{+1}\left(u+u_-\right)\\ & +\hat{P}_{ij}^{-1}\left(u+u_+\right)\hat{P}_{jk}^{0}\left(u+\Gamma\right)\hat{P}_{kl}^{+1}\left(u+u_-\right)\\ & +\hat{P}_{ij}^{+1}\left(u+u_-\right)\hat{P}_{jk}^{0}\left(u\right)\hat{P}_{kl}^{+1}\left(u+u_-\right)\\ & +\hat{P}_{ij}^{+1}\left(u+u_-\right)\hat{P}_{jk}^{0}\left(u+\Gamma\right)\hat{P}_{kl}^{+1}\left(u+u_-\right)\\ & +2\hat{P}_{ij}^{+1}\left(u+u_-\right)\hat{P}_{jk}^{+2}\left(u+2u_-\right)\hat{P}_{kl}^{+1}\left(u+u_-\right)+C.C\ ]\ \}\end{aligned} \tag{9}$$

where $u_\pm = \Gamma/2 \pm i\omega_{\mathrm{L}}$, $\mathcal{L}^{-1}$ denotes the inverse Laplace transform and $\hat{P}_{ij}^{s}(u)$ is the Laplace transform of $P_{ij}^{s}(t)$. Using standard methods of complex analysis (one can also use standard numerical or symbolic packages) we find the exact analytical expression for Mandel's Q parameter Eq. (4), based on Eqs. (9) and $\langle W\rangle$ [i.e. Eq. (6) when $T\to\infty$]. It turns out that Q is not a simple function of the model parameters; however, as we show below in certain limits simple behaviors are found.

We now consider the slow modulation limit $\nu \gg \Gamma \gg R$. Using our exact result Eq. (9) we find

$$Q \sim \frac{\xi}{2R}\left(1+\frac{e^{-2RT}-1}{2RT}\right)\frac{\left(I_+-I_-\right)^2}{\left(I_++I_-\right)}, \tag{10}$$

where $I_\pm = \Omega^2\Gamma|\frac{\Gamma}{2}+i(\omega_{\mathrm{L}}\mp\nu)|^{-2}/4$ are stationary solutions of time independent Bloch equation with frequency detuning $(\omega_{\mathrm{L}}\mp\nu)$, respectively. Similar to the line shape $\langle I\left(\omega_L\right)\rangle \sim (I_+ + I_-)/2$, Q Eq. (10) exhibits splitting with two peaks at $\omega_{\mathrm{L}} = \pm\nu$. Unlike the line shape, Q depends on measurement time T and on R. As an example we choose the parameters $\Gamma = 40$MHz, $\Omega = \Gamma/10$, $R = 1$/sec and $\omega_{\mathrm{L}} = \nu = 1$GHz, which mimics a SM coupled to a slow two level system in glass [9], we find for $RT \gg 1$, $Q = 2\times 10^5\xi$ which implies that fluctuations are very large (if compared with the fast modulation limit soon to be considered). The lengthly though straightforward derivation of Eq. (10) is based on careful analysis of the poles of $\langle W^2\rangle$ and $\langle W\rangle$ in Laplace domain. The result Eq. (10) can be easily understood in the following way. The molecule can be found in two states up + and down −, in the slow modulation limit the rate of photon emission in these two states is determined by the stationary solution of time independent Bloch equation, namely $I_\pm$, while transients are neglected. Using this physical picture one can re-derive Eq. (10) in a straightforward way. Eq. (10) describes well the experiments from Bordeaux [3] of SMS in glass, there the second order correlation function $g^{(2)}(t)$ was used to characterize the dynamics of a SM coupled to a two level system in the glass. To see the relation between these experiments and our work we recall the relation [16,19]

$$Q = \langle I\left(\omega_L\right)\rangle\left[\frac{2}{T}\int_0^T \mathrm{d}t_1\int_0^{t_1}\mathrm{d}t_2 g^{(2)}(t_2) - T\right],$$

and for slow process $g^{(2)}(t) = 1 + C\exp(-2Rt)$ with $C = (I_+ - I_-)^2/(I_+ + I_-)^2$, this expression being equivalent to Eq. (10). The question now remains whether one can go beyond this slow modulation limit and investigate faster processes based on SMS.

We therefore consider fast modulation limit for which only the $T\to\infty$ limit is relevant. Mathematically different definitions of a fast modulation exist depending on the $R\to\infty$ limiting procedure used. First consider the case $R\to\infty$ keeping ν and Γ fixed. This fast limit gives $Q = 0$. This is expected since in this case the bath is so fast the molecule cannot respond to it (e.g., the line width of the molecule is Γ). A more physically interesting case is to let $R \to \infty$ but keep the ratio $\Gamma_{\mathrm{eff}}/\Gamma \equiv \nu^2/(R\Gamma)$ finite. In this fast modulation limit the well known line shape is Lorentzian with a width $\Gamma + \Gamma_{\mathrm{eff}}$ and as mentioned it exhibits motional narrowing. Using our exact results we find in the same limit and for $\omega_{\mathrm{L}} = 0$ (i.e. at resonance)

$$Q = \frac{\xi 4\,\Omega^2\,\Gamma_{\mathrm{eff}}^2}{\Gamma\left(\Gamma + \Gamma_{\mathrm{eff}}\right)^2\,\left(\Gamma + 2\,\Gamma_{\mathrm{eff}}\right)}. \tag{11}$$

To estimate the magnitude of these fluctuations we note that the maximum of Q is found when $\Gamma_{\mathrm{eff}} = \Gamma(1+\sqrt{5})/2$ and then $Q \simeq 0.36\xi\Omega^2/\Gamma^2$. Even if we take $|\Omega|/\Gamma = 0.1$ and $\xi = 5 \times 10^{-2}$ as a reasonable estimate for weak laser field and detection efficiency, we find a small value $Q \simeq 1.8\times10^{-4}$. However values of $|Q| \le 5\times10^{-4}$ were reported in [16,22], so measurement in the fast modulation regime might also be possible.

The fast modulation limit can be analyzed using a method developed by Loring and Mukamel [13] in the context of four wave mixing. Briefly, in this limit we use a factorization approximation $\mathrm{C}_3\left(\tau_1,\tau_2,\tau_3\right) \simeq \mathrm{C}_1\left(\tau_1\right)\mathrm{C}_1\left(\tau_2\right)\mathrm{C}_1\left(\tau_3\right)$, and $\mathrm{C}_1\left(\tau\right)$ is calculated based on standard line shape theory (details will be given elsewhere [23]). This approximation works well since correlation between the state of the molecule (i.e., $+$ or $-$) during one pulse interval [say $S(t) = 1$] with that of the following pulse interval [say $S(t^{'}) = 0$] are unimportant, because the bath is fast. Employing this approach, or using Eq. (9) directly, we find

$$Q \simeq \begin{cases} \frac{\xi\Omega^2\Gamma_{\mathrm{eff}}}{2\Gamma\left(\omega_{\mathrm{L}}^2+\Gamma_{\mathrm{eff}}^2/4\right)} & \Gamma \ll \Gamma_{\mathrm{eff}} \\ \\ \frac{\xi\Omega^2\Gamma_{\mathrm{eff}}\Gamma\omega_{\mathrm{L}}^2}{\left(\omega_{\mathrm{L}}^2+\Gamma^2/4\right)^3} & \Gamma_{\mathrm{eff}} \ll \Gamma. \end{cases} \tag{12}$$

Eq. (12) exhibits several interesting behaviors **(i)** when $\Gamma \ll \Gamma_{\mathrm{eff}}$ a simple relation between the line shape and Q holds, $Q = 2\langle I\left(\omega_{\mathrm{L}}\right)\rangle/\Gamma$, and both exhibit motional narrowing, **(ii)** when $\Gamma_{\mathrm{eff}} \ll \Gamma$, $Q \sim 1/R$, this behavior showing that as the bath becomes faster the deviations from Poisson statistics become smaller (similar behavior is observed in the slow modulation limit) **(iii)** a dip in Q is found when $\Gamma_{\mathrm{eff}} \ll \Gamma$ (note that we have neglected $O(\Gamma_{\mathrm{eff}}/\Gamma)^2$ terms which give corrections at $\omega_{\mathrm{L}} = 0$).

To conclude, the three time correlation function describes line shape fluctuations for a single molecule undergoing a stationary spectral diffusion process. We know that similar correlation functions yield important information in non–linear spectroscopy [13] (i.e., beyond linear spectroscopy), similarly "Q SMS" yields detailed information on dynamical processes in the condensed phase (i.e., beyond the standard line shape theories). Depending on the bath time scale, different time orderings of the three time correlation function contribute to the line shape fluctuations and in the fast modulation limit all time orderings contribute. Our exact results for a simple Kubo-Anderson process are applicable to the experimental situation, in the slow and possibly in the intermediate modulation regime (where Q is large). Our results in the more challenging fast modulation case give the theoretical limitations on measurement accuracy needed to detect Q.

Acknowledgment This work was supported by the NSF. EB thanks M. Orrit and G. Zumofen for discussions.

---

[1] W. E. Moerner and M. Orrit, *Science* **283** 1670 (1999).

[2] W. P. Ambrose, T. Basche, and W. E. Moerner, *J. Chem. Phys* **95** 7150 (1991).

[3] A. Zumbusch, L. Fleury, R. Brown, J. Bernard, and M. Orrit, *Phy. Rev. Lett.* **70** 3584 (1993), L. Fleury, A. Zumbusch, M. Orrit, R. Brown, and J. Bernard, *J. of Lumin.* **56** 15 (1993).

[4] P. D. Reilly and J. L. Skinner, *Phys. Rev. Lett.* **71**, 4257 (1993).

[5] G. Zumofen and J. Klafter, *Chem. Phys. Lett.* **219** 303 (1994).

[6] X. Sunney Xie *Acc. Chem. Res.* **29** 598 (1996).

[7] S. A. Empedocles, D. J. Norris and M. G. Bawendi, *Phys. Rev. Lett.* **77**, 3873 (1996).

[8] H. Bach, A. Renn, G. Zumofen, and U. P. Wild, *Phys. Rev. Lett.* **82**, 2195 (1999).

[9] A. M. Boiron, P. Tamarat, B. Lounis, R. Brown, and M. Orrit, *Chem. Phys.* **247**, 119 (1999).

[10] T. Plakhotnik and D. Wasler, *Phys. Rev. Lett.* **80**, 4064 (1998), T. Plakhotnik, *Phys. Rev. B.* **59**, 4658 (1999), T. Plakhotnik, *J. of Lumin* **83-4**, 221 (1999).

[11] E. Barkai, R. Silbey, and G. Zumofen *Phys. Rev. Lett.* **84** 5339 (2000).

[12] R. Kubo, in *Fluctuations Relaxation and Resonance in Magnetic Systems* D. ter Haar ed. Plenum (New York) 1962.

[13] S. Mukamel and R. F. Loring, *J. Opt. Soc. Am. B* **3** 595 (1986), S. Mukamel, *Principles of Non linear Optical Spectroscopy* (Oxford University Press, New York, Oxford, 1995).

[14] E. Geva and J. L. Skinner, *J. Phys. Chem. B* **101**, 8920 (1997).

[15] Th. Basche, W. E. Moerner, M. Orrit, and H. Talon, *Phys. Rev. Lett.* **69**, (1992).

[16] L. Fleury, J-M. Segura, G. Zumofen, B. Hecht, and U. P. Wild, *Phys. Rev. Lett.* **84** 1148 (2000).

[17] B. Lounis, F. Jelezko, and M. Orrit, *Phys. Rev. Lett.* **78**, 3673 (1997).

[18] Ch. Brunel, B. Lounis, Ph. Tamarat, and M. Orrit, *Phys. Rev. Lett.* **81**, 2679 (1998).

[19] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics* (Cambridge University Press, New York, 1995).

[20] A. Suárez and R. Silbey, *Chem. Phys. Lett.* **218** 445 (1994).

[21] The weights are $P^0_{if}(t) = (1/2)[1+(-1)^{\delta_{if}+1}\exp(-2Rt)]$, $P^1_{+-}(t) = P^1_{-+}(t) = R\exp(-Rt)\sin(Y_1t)/Y_1$, $P^1_{\pm\pm}(t) = \exp(-Rt)\left[\cos\left(Y_1t\right) \mp i\nu\sin\left(Y_1t\right)/Y_1\right]$, $P^{-1}_{if}(t) = \mathrm{C.C}\left[P^1_{if}\left(t\right)\right]$ and $Y_1 = \sqrt{\nu^2 - R^2}$. $P^{\pm 2}_{if}(t)$ is given by $P^{\pm 1}_{if}(t)$ when ν is replaced with 2ν.

[22] As mentioned the (room temperature) experiments in [16] find sub Poissonian statistics for short measurement times, it is still unclear if and when such results can be obtained in the presence of spectral diffusion, our results indicate that in the presence of a very fast bath spectral diffusion has no effect on photon statistics.

[23] Y. Jung, E. Barkai, and R. Silbey (to be submitted)